

Table 1. Measured mean values of variables in patients with HIV infection before and one year after the initiation of ARVT

Parameter		mean	minimum	maximum
Cholesterol	before ARVT	4.69 mmol/l	2.46 mmol/l	6.66 mmol/l
	after ARVT	6.49 mmol/l	3.76 mmol/l	16.23 mmol/l
TGL	before ARVT	1.84 mmol/l	0.47 mmol/l	8.27 mmol/l
	after ARVT	4.53 mmol/l	1.01 mmol/l	29.00 mmol/l
HDL	before ARVT	0.99 mmol/l	0.26 mmol/l	1.59 mmol/l
	after ARVT	1.29 mmol/l	0.69 mmol/l	2.91 mmol/l
LDL	before ARVT	2.81 mmol/l	0.15 mmol/l	4.92 mmol/l
	after ARVT	3.74 mmol/l	0.62 mmol/l	6.58 mmol/l
Atherosclerotic index	before ARVT	2.90	0.22	5.12
	after ARVT	3.09	0.9	5.62
Glucose	before ARVT	4.69 mmol/l	3.3 mmol/l	5.70 mmol/l
	after ARVT	5.35 mmol/l	5.20 mmol/l	6.50 mmol/l
BMI	before ARVT	21.61 kg/m ²	14.18 kg/m ²	30.10 kg/m ²
	after ARVT	24.63 kg/m ²	16.96 kg/m ²	38.02 kg/m ²